



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Notes from Essex, Massachusetts, 1921.—*Uria troille*. COMMON MURRE.—On May 18, I secured a female on the Essex River, in Essex. It appeared to be in a somewhat sickly condition and its tarsi were somewhat swollen, facts which may account in part for its presence here at this late date. Its stomach was replete with the remains of fish. This is the first definite record of this species for Essex County. The mounted specimen is now in the collection of the Boston Society of Natural History.

***Sterna caspia*.** CASPIAN TERN.—Three birds appeared on the Essex River, August 31. They were seen fishing in the river and later resting with Herring Gulls on the shore. Two others were seen on September 8 near the mouth of the river. The stomach of one secured, was kindly examined for me by the U. S. Biological Survey, and its contents found to be two codlings (*Urophycis tenuis*).

***Casmerodius egretta*.** AMERICAN EGRET.—On the salt marsh of Essex River I saw a flock of three on August 19.

***Phalacrocorax carbo*.** COMMON CORMORANT.—On September 2, flying low over the water off the mouth of the Essex River, two birds of this species were identified with a flock of twenty-five Double-crested Cormorants. They were obviously larger than the latter with white breasts, and their paler bills and gular patches were in distinct contrast to the orange color of these parts in the Double-crested species. All the birds in the flock held their bills widely open as they passed, presenting a peculiar appearance.

***Coturnicops noveboracensis*.** YELLOW RAIL.—One was started in the salt marsh on September 10.

***Limosa fedoa*.** MARBLED GODWIT.—A female was killed on the Essex River on August 31. Its stomach contents were examined for me by the U. S. Biological Survey and found to consist of 14 Nereis, 6 *Gemma purpureus* and at least four other bivalves, and a seed of the pondweed (*Potamogeton pectinatus*).

***Falco rusticolus obsoletus*.** BLACK GYRFALCON.—A fine black female was shot on the edge of the salt marsh at Essex on December 10. It had been seen several times previously and when killed was said to have been attempting to carry off a domestic hen from a farmer's flock. Its crop was greatly distended and on examination was found to be crammed with the flesh and breast feathers of a Black Duck. The specimen is now mounted in the collection of the Boston Society of Natural History. —ARTHUR B. FULLER, *Boston Soc. Nat. Hist., Boston, Mass.*

Notes from Lauderdale, Fla.—At Ft. Lauderdale, Fla., February 12, 1922, I saw a male Nonpareil (*Passerina ciris*). A day or two later I saw it again in the same general vicinity where, for the third time, I saw it, February 19. On the last occasion I flushed it, together with some Grasshopper Sparrows, from a growth of tangled grass where, evidently, these finches had taken refuge for the night. Maynard refers to this